INDEX TO VOLUME 45

Ablautus rufotibialis, ethology of, 271
Acarina, Cheyletidae, 450; Oribatei, 309; Proctophyllodidae, 327
activities in artificial nests of an allodapine bee, 263
Acuera, 229
Adult brucid beetle activity as indicated by tanglefoot traps, 200
Agrotis ipsilon, larval midgut epithelium of, 242
Aldrich, J. R., T. R. Yonke, and R. D. Oetting, Histology and morphology of the abdominal scent apparatus in three alydids, 162
Alydus, histology of nymphal scent gland reservoirs in *eurinus* and *pilosulus*, 162
Amblycerus robiniae, adult seasonal activity of, 200
Ancylotela, key to species, 106; *barberi*, 100; **knulli**, 102; **lateralis**, 94; **nelsoni**, 97; **sonorae**, 99; **subcostata**, 95
A new sexually dimorphic *Empoasca*, 125
Annotated checklist of the crab spiders of North Dakota, 311
Anthonomus grandis, laboratory oviposition rates of, 252
Anthophora occidentalis, nest-building secretions of, 208
aquatic insect trap, improved, 195
Araneae, Salticidae, 324; Theridiidae, 275; Thomisidae, 311
Aspects of the biology of Costa Rican halictine bees, III. *Sphecodes kathleenae*, a social cleptoparasite of *Dialictus umbripennis*, 529; IV. *Augochlora* (*Oxystoglossella*), 18
Atopozelus **grandis**, 392
Atyeo, Warren T., see Chong K. Park
Augochlora, *nominata*, mature larva described, 27, pupa described, 30; *pura*, larval development of, 429

Bailey, Jack C., Fowden G. Maxwell, and Johnie N. Jenkins, Seasonal fluctuations in oviposition of the boll weevil in the laboratory, 252
Barr, William F., New species of *Ancylotela* from Mexico and the United States with a key to the known species, 92; also see David E. Foster
Batra, S. W. T., Notes on the behavior and ecology of the mantispid *Climaciella brunnea occidentalis*, 334; Some properties of the nest-building secretions of *Nomia*, *Anthophora*, *Hylaeus* and other bees, 208
Bell, William J., Chris Parsons, and Edward A. Martinko, Cockroach aggregation pheronomes: Analysis of aggregation tendency and species specificity, 414
Berríos-Ortiz, A., and Richard B. Selander, Sexing immature blister beetles, 376
biological control, of the greenbug, *Schizaphis graminum*, on sorghum, 324; *Metaphidippus galathea* as a possible agent, 324
biology of *Augochlora nominata* (Hymenoptera), 18; of *Osmia glauca* and *nemoris*, 523; of *O. lignaria*, 235
Bohart, G. E., P. F. Torchio, Yasuo Maeta, and R. W. Rust, Notes on the biology of *Emphoropsis pallida*, 381
Braunsapis foveata, activities of within artificial nests, 263; food transfer in artificial nests, 373
Brower, John H., "Scaleless" and "melanic," two undescribed mutations in *Plodia interpunctella*, 421; also see Robert L. Kirkpatrick
Byers, George W., Competitive supersedure by *Monobia quadridens* in nests of *Osmia lignaria*, 235; A new genus of Mecoptera from Brazil, 340

Calkins, C. O., see J. W. Matteson
Camargo, Conceição Aparecida de, Mating of the social bee *Melipona quadrifasciata* under controlled conditions, 520
Campbell, John B., see J. D. Lambley
Carlson, Donald, Comparative value of black light and cool white lamps in attracting insects to aquatic traps, 194
Chiang, H. C., see G. Wilde
chromosome numbers, in bees, 111
clades and grades in the Alydidae, 135
Clement, C. L., see R. W. Rust
cleptoparasitism in bees, 529
Climaciella brunnea occidentalis, behavior and ecology of, 334
cockroach aggregation pheromones: Analysis of aggregation tendency and species specificity, 414
Coleoptera, 122; Bostrichidae, 238; Bruchidae, 200; Buprestidae, 92; Chrysomelidae, 259, 516; Curculionidae, 252, 434; Meloidae, 128, 376, 459
Coleosoma floridanum, behavior of, 275
Columbicheyla, 450; **macroflabellata**, 450

INDEX TO VOLUME 45

Ablautus rufotibialis, ethology of, 271
Acarina, Cheyletidae, 450; Oribatei, 309; Proctophyllodidae, 327
activities in artificial nests of an allodapine bee, 263
Acuera, 229
Adult brucid beetle activity as indicated by tanglefoot traps, 200
Agrotis ipsilon, larval midgut epithelium of, 242
Aldrich, J. R., T. R. Yonke, and R. D. Oetting, Histology and morphology of the abdominal scent apparatus in three alydids, 162
Alydus, histology of nymphal scent gland reservoirs in *eurinus* and *pilosulus*, 162
Amblycerus robiniae, adult seasonal activity of, 200
Ancylotela, key to species, 106; *barberi*, 100; **knulli**, 102; **lateralis**, 94; **nelsoni**, 97; **sonorae**, 99; **subcostata**, 95
A new sexually dimorphic *Empoasca*, 125
Annotated checklist of the crab spiders of North Dakota, 311
Anthonomus grandis, laboratory oviposition rates of, 252
Anthophora occidentalis, nest-building secretions of, 208
aquatic insect trap, improved, 195
Araneae, Salticidae, 324; Theridiidae, 275; Thomisidae, 311
Aspects of the biology of Costa Rican halictine bees, III. *Sphecodes kathleenae*, a social cleptoparasite of *Dialictus umbripennis*, 529; IV. *Augochlora* (*Oxystoglossella*), 18
Atopozelus **grandis**, 392
Atyeo, Warren T., see Chong K. Park
Augochlora, *nominata*, mature larva described, 27, pupa described, 30; *pura*, larval development of, 429

Bailey, Jack C., Fowden G. Maxwell, and Johnie N. Jenkins, Seasonal fluctuations in oviposition of the boll weevil in the laboratory, 252
Barr, William F., New species of *Ancylotela* from Mexico and the United States with a key to the known species, 92; also see David E. Foster
Batra, S. W. T., Notes on the behavior and ecology of the mantispid *Climaciella brunnea occidentalis*, 334; Some properties of the nest-building secretions of *Nomia*, *Anthophora*, *Hylaeus* and other bees, 208
Bell, William J., Chris Parsons, and Edward A. Martinko, Cockroach aggregation pheronomes: Analysis of aggregation tendency and species specificity, 414
Berríos-Ortiz, A., and Richard B. Selander, Sexing immature blister beetles, 376
biological control, of the greenbug, *Schizaphis graminum*, on sorghum, 324; *Metaphidippus galathea* as a possible agent, 324
biology of *Augochlora nominata* (Hymenoptera), 18; of *Osmia glauca* and *nemoris*, 523; of *O. lignaria*, 235
Bohart, G. E., P. F. Torchio, Yasuo Maeta, and R. W. Rust, Notes on the biology of *Emphoropsis pallida*, 381
Braunsapis foveata, activities of within artificial nests, 263; food transfer in artificial nests, 373
Brower, John H., "Scaleless" and "melanic," two undescribed mutations in *Plodia interpunctella*, 421; also see Robert L. Kirkpatrick
Byers, George W., Competitive supersedure by *Monobia quadridens* in nests of *Osmia lignaria*, 235; A new genus of Mecoptera from Brazil, 340

Calkins, C. O., see J. W. Matteson
Camargo, Conceição Aparecida de, Mating of the social bee *Melipona quadrifasciata* under controlled conditions, 520
Campbell, John B., see J. D. Lambley
Carlson, Donald, Comparative value of black light and cool white lamps in attracting insects to aquatic traps, 194
Chiang, H. C., see G. Wilde
chromosome numbers, in bees, 111
clades and grades in the Alydidae, 135
Clement, C. L., see R. W. Rust
cleptoparasitism in bees, 529
Climaciella brunnea occidentalis, behavior and ecology of, 334
cockroach aggregation pheromones: Analysis of aggregation tendency and species specificity, 414
Coleoptera, 122; Bostrichidae, 238; Bruchidae, 200; Buprestidae, 92; Chrysomelidae, 259, 516; Curculionidae, 252, 434; Meloidae, 128, 376, 459
Coleosoma floridanum, behavior of, 275
Columbicheyla, 450; **macroflabellata**, 450

comparative courtship behavior of *Negalius*, *Phodaga*, and *Cordylospasta*, three closely related genera of blister beetles, 459
comparative value of black light and cool white lamps in attracting insects to aquatic traps, 194
comparison of microwave and infrared radiation to control rice weevils in wheat, 434
competitive supersedure by *Monobia quadridens* in nests of *Osmia lignaria*, 235
control of northern and western corn rootworms, *Diabrotica longicornis* and *D. virgifera*, corn yield data, and weed control afforded by certain tillage practices, 516
Cordylospasta opaca, male courtship behavior of, 468
Costanana, 227, key to species, 491; **asymmetria**, 497; **cella**, 497; **costata**, 498; **dunda**, 229; *flavicosta*, 498; **flavina**, 496; **helvacosta**, 496; **minuta**, 499; *nana*, 492; *praecellens*, 499
courtship behavior, in male of *Meloe strigulosus*, 128; in three genera of blister beetles, 459
Culumana, 233, key to males, 405; **exeava**, 408; **fascia**, 409; **incida**, 411; **pilana**, 409; **tormina**, 412; **torqua**, 233; **turba**, 413; **vitta**, 408
Curtara, 231; **samera**, 231
Cutler, Bruce, Notes on the behavior of *Coleosoma floridanum* Banks, 275

DeLong, Dwight M., and Paul H. Freytag, Studies of the Gyponinae: A key to the known genera and descriptions of five new genera, 218; The genus *Folicana* and nine new species, 282; The genus *Culumana* and seven new species, 405; The genus *Costanana* and five new species, 491
Diabrotica, egg hatch in *longicornis* and *virgifera*, 259; control of *longicornis* and *virgifera*, 516
Dialictus umbripennis, cleptoparasite of, 529
Dickerson, Williard A., and William H. Kearby, The identification and distribution of the tip moths of the genus *Rhyacionia* in Missouri, 542
Diptera, Asilidae, 271; Ceratopogonidae, 476; Leptogastridae, 7
direct food transferring behavior in bees, 373
effect of photoperiod on the development of the sawfly *Neodiprion taedae linearis*, 1
Eickwort, George C., and Kathleen R. Eickwort, Aspects of the biology of Costa Rican halictine bees: III, *Sphecodes kathleenae*, a social cleptoparasite of *Dialictus umbripennis*, 529; IV, *Augochlora* (*Oxystoglossella*), 18
Eickwort, George C., and Gerald I. Stage, A new subgenus of Neotropical *Sphecodes* cleptoparasitic upon *Dialictus*, 500
Eickwort, Kathleen R., see George C. Eickwort
Emphoropsis pallida, biology of, 381
Empoasca **dimorpha**, 125
Enns, W. R., see S. E. Thewke
Entylia bacriana, alimentary canal of, 46
Erythroneura **amboyensis**, 431; **johnsoni**, 432; **parisensis**, 431; **rubulna**, 433; **shumiquera**, 431
ethology of *Ablautus rufotibialis* on the Pawnee grasslands IBP site, 271
Eutogenes **pinicola**, 453

Fairchild, M. L., see J. C. Reese
five new species of *Erythroneura*, 430
flights and swarms of the ant *Formica obscuripes*, 254
fine structure of the midgut epithelium in larvae of *Agrotis ipsilon*, 242
Folicana, 226; **acrita**, 289; **acuera**, 282; **cornea**, 287; **digita**, 283; **figura**, 285; **furcata**, 288; **guianae**, 287; **lata**, 283; **nota**, 226; **perula**, 285
food preferences of grasshoppers in six planted pastures in eastern Kansas, 59
Formica obscuripes, flights and swarms of, 254
Foster, David E., and William F. Barr, Notes on the distribution and bionomics of some North American Cleridae, 122
Freytag, Paul H., see Dwight M. DeLong

genital morphology and species of the Eastern Hemisphere genus *Lobus*, 7
genus *Nyctiophylax* in North America, 172
Godfrey, George L., The last larval instar of *Lithacodia albidula*, 427
grasshoppers, food preferences of, 59
Gundu Rao, H. R., and Donald A.

Wilbur, Loss of wheat weight from feeding of lesser grain borer, 238
Gyponinae, key to genera, 218; studies of, 282, 405, 491

Hamilton, K. G. Andrew, The insect wing: Part II, Vein homology and the archetypal insect wing, 54; Part III, Venation of the orders, 145; Part IV, Venational trends and the phylogeny of the winged orders, 295
Hart, Elwood R., A new species of the genus *Atopozelus*, with a review of the genus, 392
Hemiptera, Alydidae, 135, 162; Reduviidae, 392
Hepner, Leon W., Five new species of *Erythroneura*, 431
Hibbs, E. T., see G. Wilde
histology and morphology of the abdominal scent apparatus in three alydids, 162
Homoptera, Cicadellidae, 125, 218, 282, 405, 430, 491; Membracidae, 46
Horner, Norman V., *Metaphidippus galathea* as a possible biological control agent, 324
host records for some North American Pompilidae, Second supplement, 181
Howe, W. L., see J. W. Matteson
Hylaeus cressoni, nest-building secretions of, 208
Hymenoptera, Anthophoridae, 111, 381; Apidae, 111, 373, 520; Apoidea, 208, 439; Diprionidae, 1; Eumenidae, 235; Formicidae, 254; Halictidae, 18, 111, 500, 529; Megachilidae, 51, 235, 523; Pompilidae, 181; Sphecidae, 397; Xylocopinae, 263

identification and distribution of the tip moths of the genus *Rhyacionia* in Missouri, 542
insect wing, vein homology and the archetypal insect wing, 54; venation of the orders, 145; venational trends and the phylogeny of the winged orders, 295
Issikiella, 340; **pulchra**, 342

Jalil, M., A note on the life cycle of *Platynothrus peltifer*, 309
Jenkins, Johnie N., see Jack C. Bailey

Kearby, William H., see William A. Dickerson
Kerr, Warwick Estevam, Numbers of chromosomes in some species of bees, 111
Kirkpatrick, Robert L., John H. Brower, and Elvin W. Tilton, A comparison of microwave and infrared radiation to control rice weevils, 434
Knutson, Herbert, see J. D. Lambley
Kopp, Dennis D., and Thomas R. Yonke, Morphological observations on the alimentary canal of *Entylia bactriana*, 46
Kurczewski, Edmund J., see Frank E. Kurczewski
Kurczewski, Frank E., Observations on the nesting behavior of *Oxybelus cressonii* and *O. exclamans*, 397; and Edmund J. Kurczewski, Host records for some North American Pompilidae, Second supplement, 181

Lambley, J. D., John B. Campbell, and Herbert Knutson, Food preferences of grasshoppers in six planted pastures in eastern Kansas, 59
Last larval instar of *Lithacodia albidula*, 427
Lavigne, Robert, Ethology of *Ablautus rufotibialis* on the Pawnee grasslands IBP site, 271
Lawson, D. E., see G. Wilde
Lepidoptera, Noctuidae, 242, 427; Olethreutidae, 542; Phycitidae, 421
light traps, comparative value of black and cool white lamp, 194
Lithacodia albidula, last larval instar of, 427
Lobus, 7, key to species, 10; *Lobus* (**Lobus**), 11, **janssensi**, 11, **kenyae**, 12, **keralae**, 13, **liberiae**, 14, **unilineatus**, 14; *Lobus* (**Nanilobus**), 15, **himalayae**, 15, **guineae**, 16
Loss of wheat weight from feeding of lesser grain borer, 238

Maeta, Yasuo, see G. E. Bohart
Martin, Charles H., Genital morphology and species of the Eastern Hemisphere genus *Lobus*, 7
Martinko, Edward A., see William J. Bell
McKnight, M. E., and A. D. Tagestad, *Megachile centuncularis* nest in carpenterworm gallery, 51
Mathwig, John E., Adult brucid beetle activity as indicated by tanglefoot traps, 200
Mating of the social bee *Melipona quadrifasciata* under controlled conditions, 520

Matteson, J. W., C. O. Calkins, and W. L. Howe, Control of northern and western corn rootworms, *Diabrotica longicornis* and *D. virgifera*, corn yield data, and weed control afforded by certain tillage practices, 516
Maxwell, Fowden G., see Jack C. Bailey
May, Dorothy G. K., Water uptake during larval development of a sweat bee, *Augochlora pura*, 439
Mecoptera, Bittacidae, 340
Megachile centuncularis nest in carpenterworm gallery, 51
Megalotomus, histology of nymphal scent gland reservoirs in *M. quinquespinosus*, 162
"melanic" mutation in *Plodia interpunctella*, 421
Meloe (*Meloe*) *strigulosus*, male courtship behavior of, 128
Metaphidippus galathea as a possible biological control agent, 324
Michener, Charles D., Activities within artificial nests of an allodapine bee, 263; Direct food transferring behavior in bees, 373
Monobia quadridens, nest building of, 235
morphology, of the alimentary canal of *Entylia bactriana*, 46; genital, of the genus *Lobus*, 7
Morse, John C., The genus *Nyctiophylax* in North America, 172

Negalius marmoratus, male courtship behavior of, 462
Neivamyrmex, key, 348; **texanus**, 353
Neobezzia, a new Neotropical biting midge genus of the tribe Sphaeromiini, 476; **Neobezzia**, 477, key to species (females), 478; **albitarsis**, 478, **amnicola**, 480, **blantoni**, 482, **brasiliae**, 484, **clavipes**, 485, **costaricae**, 486, **fittkaui**, 489
Neodiprion taedae linearis, photoperiod and larval development, 1
Neoeucheyla **beeri**, 455
nest-building secretions in bees, 208
nesting behavior, in wasps, 397
Neuroptera, Mantispidae, 334
new genus, and three new species of cheyletid mites from Missouri and Michigan, 450; of allodectine feather mites from hummingbirds, 327; of Mecoptera from Brazil, 340
new species, of *Ancylotela* from Mexico and the United States with a key to the known species, 92; of the genus *Atopozelus*, with a review of the genus, 392
new subgenus of Neotropical *Sphecodes* cleptoparasitic upon *Dialictus*, 500
Nomia melanderi, nest-building secretions of, 208
note on the life cycle of *Platynothrus peltifer*, 309
notes, on the behavior and ecology of the mantispid *Climaciella brunnea occidentalis*, 334; on the behavior of *Coleosoma floridanum*, 275; on the biology of *Emphoropsis pallida*, 381; on the distribution and bionomics of some North American Cleridae, 122
numbers of chromosomes in some species of bees, 111
Nyctiophylax, key to males, 178; **banksi**, 173; **denningi**, 175

observations on the nesting behavior of *Oxybelus cressonii* and *O. exclamans*, 397
Oetting, R. D., see J. R. Aldrich
Orthoptera, Acrididae, 59; Blattidae, 414
Osmia, *lignaria*, nest supersedure by eumenid wasp, 235; biology of *O. glauca*, 524; of *O. nemoris*, 527

Park, Chong K., and Warren T. Atyeo, A new genus of allodectine feather mites from hummingbirds, 327
Parsons, Chris, see William J. Bell
pheromones, aggregation, effect of on resting site of cockroaches, 405
Phodaga alticeps, male courtship behavior of, 467
photoperiod, effect of on larval development, 1
Pinto, John D., The sexual behavior of *Meloe* (*Meloe*) *strigulosus*, 128; Comparative courtship behavior of *Negalius*, *Phodaga*, and *Cordylospasta*, three closely related genera of blister beetles, 459
Platynothrus peltifer, life cycle of, 309
Plodia interpunctella, mutations in, 421
radiation, microwave and infrared, as rice weevil control, 434
Ratanaworabhan, Niphan Chanthawanich, see Willis W. Wirth
Reese, J. C., T. R. Yonke, and M. L. Fairchild, Fine structure of the midgut epithelium in larvae of *Agrotis ipsilon*, 242
reviews, Ross and Ricker, The classifi-

cation, evolution, and dispersal of the winter stonefly genus *Allocapnia*, 142; Schneirla, Army ants, 141; Shor, Fossils and flies, 268
Rhizopertha dominica, weight loss in wheat fed upon by, 238
Rhyacionia, key to pupae, 546; distribution and identification of *frustrana*, *buoliana*, *rigidana*, 542
Romero S., Jose I., see Robert F. Ruppel
Ruppel, Robert F., and Jose I. Romero S., A new sexually dimorphic *Empoasca*, 125
Rust, R. W., see G. E. Bohart, and S. L. Clement, The biology of *Osmia glauca* and *O. nemoris*, 523

Sauer, Richard J., An annotated checklist of the crab spiders of North Dakota, 311
"Scaleless" and "melanic," two undescribed mutations in *Plodia interpunctella*, 421
scent gland-reservoirs, abdominal, histology and morphology of in alydids, 162
Schaefer, C. W., Clades and grades in the Alydidae, 135
Schizodectes, 329, **hiterminalis**, 333
seasonal fluctuations in oviposition of the boll weevil in the laboratory, 252
Selander, Richard B., see A. Berrios-Ortiz
Sexing immature blister beetles, 376
sexual, behavior of *Meloe* (*Meloe*) *strigulosus*, 128; dimorphism, internal and external, in immature blister beetles, 376
Sitophilus oryzae, the rice weevil, control of, 434
some properties of the nest-building secretions of *Nomia*, *Anthophora*, *Hylaeus* and other bees, 208
Sphecodes (**Microsphecodes**), 501; (*M.*) **kathleenae**, 502; *kathleenae* as a social cleptoparasite, 529; (*M.*) **dominicanus**, 509
Stage, Gerald I., see George C. Eickwort
studies of the Gyponinae: A key to the known genera and descriptions of five new genera, 218; The genus *Folicana* and nine new species, 282; The genus *Culumana* and seven new species, 405; The genus *Costanana* and five new species, 491

Tagestad, A. D., see M. E. McKnight
Talbot, Mary, Flights and swarms of the ant *Formica obscuripes*, 254
taxonomy of *Neivamyrmex texanus*, n. sp., *N. nigrescens* and *N. californicus*, with distribution maps and keys to the species of *Neivamyrmex* of the United States, 347
Thewke, S. E., and W. R. Enns, A new genus and three new species of cheyletid mites from Missouri and Michigan, 450
tillage practices, ineffective in control of northern and western corn rootworms, 516
Tilton, Elvin W., see Robert L. Kirkpatrick
Torchio, P. F., see G. E. Bohart
Tricoptera, Polycentropidae, 172

variation in egg hatch among western and northern corn rootworms from midwest, 259
vein homology, of insect wing, 54
venation of insect orders, 145, 295

Warren, L. O., see Melvin Wingfield
water uptake during larval development of a sweat bee, *Augochlora pura*, 439
Watkins, Julian F., II, The taxonomy of *Neivamyrmex texanus*, n. sp., *N. nigrescens* and *N. californicus*, with distribution maps and keys to the species of *Neivamyrmex* of the United States, 347
Wilde, G., H. C. Chiang, E. T. Hibbs, and D. E. Lawson, Variation in egg hatch among western and northern corn rootworms from six midwestern States, 259
wing, vein homology of, 54; venation of the orders, 145; venational trends and phylogeny of winged orders, 295
Wingfield, Melvin, and L. O. Warren, The effect of photoperiod on the development of the sawfly *Neodiprion taedae linearis*, 1
Wirth, Willis W., and Niphan Chanthawanick Ratanaworabhan, *Neobezzia*, a new Neotropical biting midge genus of the tribe Sphaeromiini, 476

Yonke, T. R., see J. R. Aldrich, see Dennis D. Kopp, see J. C. Reese